



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

I. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.; CHARLES C. CROSS, Whaleyville, Va.; and H. C. WHITAKER, Ph. D., Professor of Mathematics, Manual Training School, Philadelphia, Pa.

Take $BC=a$, $AC=b$, and $AB=c$, and put the medians equal, respectively, m_a , m_b , and m_c . Then

$$m_a = \frac{1}{2}\sqrt{2(b^2 + c^2) - a^2},$$

$$m_b = \frac{1}{2}\sqrt{2(a^2 + c^2) - b^2},$$

$$m_c = \frac{1}{2}\sqrt{2(a^2 + b^2) - c^2}.$$

But the medians of a triangle intersect at a common point two-thirds of the distance from the vertex to the middle of the opposite side. Whence

$$GA^2 = (\frac{2}{3}m_a)^2 = \frac{2}{3}b^2 + \frac{2}{3}c^2 - \frac{1}{3}a^2,$$

$$GB^2 = (\frac{2}{3}m_b)^2 = \frac{2}{3}a^2 + \frac{2}{3}c^2 - \frac{1}{3}b^2,$$

$$GC^2 = (\frac{2}{3}m_c)^2 = \frac{2}{3}a^2 + \frac{2}{3}b^2 - \frac{1}{3}c^2.$$

$$\therefore GA^2 + GB^2 + GC^2 = \frac{1}{3}(a^2 + b^2 + c^2), \text{ and } 3(GA^2 + GB^2 + GC^2) = a^2 + b^2 + c^2 = BC^2 + AC^2 + AB^2.$$

II. Solution by J. W. YOUNG, Fellow and Assistant, Ohio State University, Columbus, O.; P. S. BERG, B. Sc., Principal of Schools, Larimore, N. D.; and J. SCHEFFER, A. M., Hagerstown, Md.

Let ABC be the triangle, AM , BN , and CL the medians, and G the intersection of the medians. Then by a well-known theorem,

$$AC^2 + CB^2 = 2CL^2 + 2AL^2 = 2CL^2 + \frac{1}{2}AB^2.$$

Similarly,

$$CB^2 + BA^2 = 2BN^2 + \frac{1}{2}AB^2,$$

$$BA^2 + AC^2 = 2AM^2 + \frac{1}{2}CB^2.$$

Adding and dividing by 2, we have

$$AC^2 + CB^2 + BA^2 = CL^2 + BN^2 + AM^2 + \frac{1}{2}(AB^2 + AC^2 + CB^2),$$

$$\text{or } AC^2 + CB^2 + BA^2 = \frac{4}{3}(CL^2 + BN^2 + AM^2),$$

(since G divides medians in the ratio 2:1)

$$= \frac{4}{3}(\frac{1}{4}CG^2 + \frac{1}{4}BG^2 + \frac{1}{4}AG^2) = 3(CG^2 + BG^2 + AG^2).$$

The same propositions can very easily be proven analytically.

Solved in a similar manner by COOPER D. SCHMITT, G. B. M. ZERR, WALTER H. DRANE, and CHAS. C. CROSS.

124. Proposed by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics in Drury College, Springfield, Mo.

Every conic that passes through all the foci of a conic is a rectangular hyperbola. [From Charlotte A. Scott's *Modern Analytical Geometry*.]

I. Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, O.

If $u=0$(1) be the general equation of the second degree, we have the well-known equations

$$Fx+Gy-H=Cxy\ldots\ldots(2),$$

$$2Gx-2Fy-A+B=C(x^2-y^2)\ldots\ldots(3),$$

for the foci, A , B , etc., being, as usual, the various minors of

$$\begin{vmatrix} a, & h, & g \\ h, & b, & f \\ g, & f, & c \end{vmatrix}$$

Either (2) or (3) is the rectangular hyperbola.

II. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

"If through each of the two imaginary points at infinity on any circle two tangents be drawn to the conic, these tangents will form a quadrilateral whose vertices will be the real and imaginary foci of the conic."

Both Puckle and Salmon solve this problem. Salmon's proof expanded is as follows :

Find the condition that $x-x'+(y-y')\sqrt{-1}=0$ should touch the conic

$$ax^2+2hxy+by^2+2gx+2fy+c=0.$$

The condition that $mx+ny+p=0$ touches the same conic is

$$(bc-f^2)m^2+(ca-g^2)n^2+(ab-h^2)p^2+2(gh-af)pn+2(hf-bg)pm+2(fg-ch)mn=0,$$

$$\text{or } Am^2+Bn^2+Cp^2+2Fpn+2Gpm+2Hmn=0.$$

Now $m=1$, $n=\sqrt{-1}$, $p=-x'-y'\sqrt{-1}$ or dropping the accents $p=-(x+y\sqrt{-1})$.

$$\therefore A-B+Cx^2-Cy^2+2Cxy\sqrt{-1}-2Fx\sqrt{-1}+2Fy-2Gx-2Gy\sqrt{-1}+2H\sqrt{-1}=0.$$

$$\therefore C(x^2-y^2)+2Fy-2Gx+A-B=0\ldots\ldots(1),$$

$$\text{and } 2Cxy-2Fx-2Gy+2H=0, \text{ or}$$

$$Cxy-Fx-Gy+H=0\ldots\ldots(2).$$

The intersections of (1) and (2), both of which are rectangular hyperbolas, determine the foci.